



# HUA YI SECONDARY SCHOOL 4-G3 / PRELIMINARY EXAM 2024 5-G2

NAME

CLASS

--	--

INDEX  
NUMBER

--	--

## ADDITIONAL MATHEMATICS PAPER 1

4049/01

22 September 2024

2 hour 15 minutes

Candidates answer on the Question Paper  
No Additional Materials is required.

### READ THESE INSTRUCTIONS FIRST

Write your Name, Class, and Index Number in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue, or correction fluid.

Answer **all** the questions.

The number of marks is given in the brackets [ ] at the end of each question or part question.

If working is needed for any question it must be shown with the answer. Omission of essential working will result in loss of marks.

The total number of marks for this paper is 90.

The use of an approved scientific calculator is expected, where appropriate. If the degree of accuracy is not specified in the question and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place. For  $\pi$ , use either your calculator value or 3.142.

This document consists of **21** printed pages and **1** blank page.

© HYSS 2024

No part of this document may be reproduced in any form or transmitted in any form or by any means without the prior permission of Hua Yi Secondary School.

[Turn Over

*Mathematical Formulae***1. ALGEBRA***Quadratic Equation*For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial expansion*

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$ **2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for  $\triangle ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$

- 1 A cuboid has a base area of  $(7 + 4\sqrt{5})\text{cm}^2$  and a volume of  $(16 + 18\sqrt{5})\text{cm}^3$ . Find, without using a calculator, the height of the cuboid, in cm, in the form  $(a + b\sqrt{5})$ , where  $a$  and  $b$  are integers.

[3]

- 2 The curve  $5x - xy = 20$  and the line  $x - 2y - 3 = 0$  intersects at the points  $A$  and  $B$ .  
Find the  $y$ -coordinate of  $A$  and of  $B$ .

[3]

- 3 (a) Express  $12x - 13 - 3x^2$  in the form  $a(x + b)^2 + c$  and hence state the coordinates of the turning point of the curve  $y = 12x - 13 - 3x^2$ . [4]

- (b) State the range of  $k$  such that  $y = k$  will intersect the curve  $y = 12x - 13 - 3x^2$ . [1]

4 Integrate  $\frac{4}{5x+1} - \frac{6}{x^3}$  with respect to  $x$ .

[2]

5 Express  $\frac{7x^2-17x+1}{(x^2+1)(2-3x)}$  in partial fractions.

[5]

- 6 Given that  $x^5 + ax^3 + bx^2 - 3 \equiv (x^2 - 1)Q(x) - x - 2$ , where  $Q(x)$  is a polynomial,

(a) state the degree of  $Q(x)$ ,

[1]


(b) show that  $a = -2$  and  $b = 1$ ,

[3]

(c) find the polynomial  $Q(x)$ .

[3]

7



The sketch above shows part of the graph of  $y = a \sin\left(\frac{x}{b}\right) + c$ , where  $x$  is in degrees.

(a) Explain why  $c = 1$ .

[2]

(b) State the value of  $b$ .


[1]

(c) Find the value of  $m$  and explain how you get it.

[2]




- 8 The figure shows the curve  $y = x^2$  and the point  $R(1,0)$ . The variable point  $P(p, 0)$  moves along the  $x$  axis and  $PQ$  is vertical. It is given that  $p$  decreases at the rate of 1.2 units per second.



- (a) Show that the area of the triangle  $PQR$ ,  $A$  units<sup>2</sup>, is  $A = \frac{1}{2}p^2 - \frac{1}{2}p^3$ . [2]

- (b) Find the rate at which  $A$  is increasing at the instant when  $p = -7$ . [4]

- 9 From a rectangular piece of metal of width 2m and length 6m, two squares of side  $x$  m and two rectangles of sides  $x$  m and  $(x + y)$  m are removed as shown. The metal is then folded about the dotted lines to give a closed box with height  $x$  m.



- (a) Show that the volume of the box,  $V \text{ m}^3$ , is given by  $V = 2x^3 - 8x^2 + 6x$ .

[3]


(b) Given that  $x$  can vary, find the stationary value of  $V$ .

[4]

(c) Show that this value of  $V$  is a maximum.

[2]

10 The diagram shows a triangle  $ABC$  with vertices at  $A(0, 6)$ ,  $B(8, 15)$  and  $C(20, k)$ .



(a) Given that  $AB = BC$ , find the value of  $k$ .

[4]

(b)  $D$  is a point on the  $x$  axis such that  $ABCD$  is a kite. Find the coordinates of  $D$ .

[4]

(c) Hence find the area of the kite  $ABCD$ .

[2]

11 (a) Prove the identity  $\cot \theta + \frac{\sin \theta}{1 + \cos \theta} = \operatorname{cosec} \theta$ .

[3]

(b) Hence solve the equation  $\cot 3\theta + \frac{\sin 3\theta}{1+\cos 3\theta} = -2$  for  $-90^\circ \leq \theta \leq 90^\circ$ .

[4]

12 (a) Solve  $6^{2x-1} = 4^x \times 5$ .

[4]

(b) Given that  $\log_x 3 = p$ , express  $\log_3 \frac{27}{x}$  in terms of  $p$ .


[2]



(c) Solve the equation  $\lg x - 1 = \lg(x - 1)$

[3]

13 The diagram shows part of the curve  $y = 8 - e^{-3x}$  which crosses the axes at  $A$  and at  $B$ .



(a) Show that the  $x$  coordinate of  $A$  is  $-\ln 2$ .

[3]


(b) The normal to the curve at  $B$  meets the  $x$  axis at  $C$ . Find the coordinates of  $C$ .

[4]

(c) Find the area of the shaded region.

[4]

14



The diagram shows a circle, centre  $O$ , with diameter  $AB$ . The point  $C$  lies on the circle. The tangent to the circle at  $A$  meets  $BC$  extended at  $D$ . The tangent to the circle at  $C$  meets the line  $AD$  at  $E$ .

- (a) Prove that triangles  $EOA$  and  $EOC$  are congruent.

[3]

(b) Prove that triangles  $ADC$  and  $BAC$  are similar.

[3]

(c) Hence prove that  $AC^2 = CD \times BC$ .

[2]

**End of Paper**

**BLANK PAGE**

- 1 A cuboid has a base area of  $(7 + 4\sqrt{5})\text{cm}^2$  and a volume of  $(16 + 18\sqrt{5})\text{cm}^3$ . Find, without using a calculator, the height of the cuboid, in cm, in the form  $(a + b\sqrt{5})$ , where  $a$  and  $b$  are integers.

[3]

$$\begin{aligned} h &= \frac{16+18\sqrt{5}}{7+4\sqrt{5}} \times \frac{7-4\sqrt{5}}{7-4\sqrt{5}} \\ &= \frac{112-64\sqrt{5}+126\sqrt{5}-360}{-31} \\ &= 8 - 2\sqrt{5} \end{aligned}$$

- 2 The curve  $5x - xy = 20$  and the line  $x - 2y - 3 = 0$  intersects at the points  $A$  and  $B$ . Find the  $y$ -coordinate of  $A$  and of  $B$ .

[3]

*sub  $x = 2y + 3$  into first equation,*

$$5(2y + 3) - y(2y + 3) = 20$$

$$2y^2 - 7y + 5 = 0$$

$$(2y - 5)(y - 1) = 0 \text{ or by quadratic formula}$$

$$y = 2.5 \text{ or } 1$$



- 3 (a) Express  $12x - 13 - 3x^2$  in the form  $a(x + b)^2 + c$  and hence state the coordinates of the turning point of the curve  $y = 12x - 13 - 3x^2$ . [4]

$$\begin{aligned}
 & -3(x^2 - 4x) - 13 && \text{or} && -3\left(x^2 - 4x + \frac{13}{3}\right) \text{-----} \\
 & = -3[(x - 2)^2 - 4] - 13 && \text{or} && = -3\left[(x - 2)^2 - 4 + \frac{13}{3}\right] \\
 & = -3(x - 2)^2 - 1
 \end{aligned}$$

Turning point (2, -1)

- (b) State the range of  $k$  such that  $y = k$  will intersect the curve  $y = 12x - 13 - 3x^2$ . [1]

$$k \leq -1$$

- 4 Integrate  $\frac{4}{5x+1} - \frac{6}{x^3}$  with respect to  $x$ .

[2]

$$\int \frac{4}{5x+1} - 6x^{-3} dx = \frac{4 \ln(5x+1)}{5} + \frac{3}{x^2} + c$$

- 5 Express  $\frac{7x^2-17x+1}{(x^2+1)(2-3x)}$  in partial fractions.

[5]

$$\frac{Ax+B}{x^2+1} + \frac{C}{2-3x} \quad \text{M1}$$

$$7x^2 - 17x + 1 = (Ax + B)(2 - 3x) + C(x^2 + 1)$$

$$A = -4,$$

$$B = 3$$

$$C = -5$$

$$\frac{3-4x}{x^2+1} - \frac{5}{2-3x}$$

- 6 Given that  $x^5 + ax^3 + bx^2 - 3 \equiv (x^2 - 1)Q(x) - x - 2$ , where  $Q(x)$  is a polynomial.

(a) State the degree of  $Q(x)$ .

[1]

$$\text{Degree of } Q(x) = 3$$

(b) Show that  $a = -2$  and  $b = 1$ .

[3]

$$\text{Sub } x = 1, \quad a + b = -1 \text{ -----equation}$$

$$\text{Sub } x = -1, \quad -a + b = 3 \text{ -----equation}$$

Solve simultaneous equations,

$$a = -2 \text{ and } b = 1 \text{ -----}$$

(c) Find the polynomial  $Q(x)$ .

[3]


$$Q(x)(x^2 - 1) = x^5 - 2x^3 + x^2 - 3 + x + 2$$

$$Q(x) = (x^5 - 2x^3 + x^2 - 3 + x + 2) \div (x^2 - 1)---$$

By long division or comparing terms,

$$Q(x) = x^3 - x + 1$$

7



The sketch above shows part of the graph of  $y = a \sin\left(\frac{x}{b}\right) + c$ , where  $x$  is in degrees.

- (a) Explain why  $c = 1$ .

[2]

amplitude is 3.  
So the maximum value of  $y$  is 4,  $3+c=4$ , means that  $c = 1$ . for showing how to get  $c$

- (b) State the value of  $b$ .

[1]

$$b = 2.$$


- (c) Find the value of  $m$  and explain how you get it.

[2]

The period is  $720^\circ$ . -----

$$\text{Hence } m = 180 + 720 = 900$$

- 8 The figure shows the curve  $y = x^2$  and the point  $R(1,0)$ . The variable point  $P(p, 0)$  moves along the  $x$  axis and  $PQ$  is vertical.  $p$  is decreasing at the rate of 1.2 units per second.



- (a) Show that the area of the triangle  $PQR$ ,  $A$  units<sup>2</sup>, is  $A = \frac{1}{2}p^2 - \frac{1}{2}p^3$ . [2]

getting  $y$  coordinate of  $Q = p^2$  -----

$$\begin{aligned} A &= \frac{1}{2}(1-p)(p^2) \\ &= \frac{1}{2}p^2 - \frac{1}{2}p^3 \text{ (Shown)} \end{aligned}$$

- (b) Find the rate at which  $A$  is increasing at the instant when  $p = -7$  units. [4]

$$\frac{dA}{dt} = \frac{dA}{dp} \times \frac{dp}{dt}$$


$$\frac{dA}{dt} = p - \frac{3}{2}p^2 \text{-----}$$

$$\frac{dA}{dt} = \frac{dA}{dp} \times \frac{dp}{dt} \text{-----}$$

When  $p = -7$ ,

$$\begin{aligned} \frac{dA}{dt} &= -80.5 \times -1.2 \\ &= 96.6 \text{ unit}^2 \text{ per second} \end{aligned}$$

- 9 From a rectangular piece of metal of width 2m and length 6m, two squares of side  $x$  m and two rectangles of sides  $x$  m and  $(x + y)$  m are removed as shown. The metal is then folded about the dotted lines. To give a closed box with height  $x$  m.



- (a) Show that the volume of the box,  $V \text{ m}^3$ , is given by  $V = 2x^3 - 8x^2 + 6x$ . [3]

$$\text{Length} = y, \text{ breadth} = 2 - 2x, \text{ height} = x$$

$$x + y + x + y = 6 \rightarrow y = 3 - x$$

$$\begin{aligned} V &= xy(2 - 2x) \\ &= x(3 - x)(2 - 2x) \\ &= 2x^3 - 8x^2 + 6x \text{ (shown) } \end{aligned}$$

- (b) Given that  $x$  can vary, find the stationary value of  $V$ .

[4]

$$\frac{dv}{dx} = 6x^2 - 16x + 6$$

$$\frac{dv}{dx} = 0 \rightarrow 6x^2 - 16x + 6 = 0$$

Solve using quadratic formula ( you need to show working)

$$x = 2.215 \text{ (reject) or } 0.4514$$


Stationary value of  $V = 1.26 \text{ m}^3$

- (c) Shows that this value of  $V$  is the maximum.

[2]

Show either by first or second derivative test

10 The diagram shows a triangle  $ABC$  with vertices at  $A(0, 6)$ ,  $B(8, 15)$  and  $C(20, k)$ .



- (a) Given that  $AB = BC$ , find the value of  $k$ . [4]

$$\text{Since } AB = BC, \text{ it means } 8^2 + 9^2 = 12^2 + (15 - k)^2 \text{ ----}$$

$$(15 - k)^2 = 1$$

$$(15 - k) = 1 \text{ or } (k - 15) = -1$$

$$k = 14 \text{ ----}$$



- (b)  $D$  is a point on the  $x$  axis such that  $ABCD$  is a kite. Find the coordinates of  $D$ . [4]

$$M_{AC} = \frac{2}{5}, \text{ so } M_{BD} = -\frac{5}{2}$$

Midpoint of  $AC = (10, 10)$  -----

$$\text{Find Equation of } BD : y = -\frac{5}{2}x + 35$$

Get  $D(14, 0)$

- (c) Hence, find the area of the kite  $ABCD$ . [2]

Shoelace method or sum of area of triangles -----

$$\text{Area of kite } ABCD = 174 \text{ units}^2 \text{-----}$$

11 (a) Prove the identity  $\cot \theta + \frac{\sin \theta}{1+\cos \theta} = \operatorname{cosec} \theta$ .

[3]

$$\begin{aligned} LHS &= \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{1+\cos \theta} \\ &= \frac{\cos \theta + \frac{\sin^2 \theta}{\cos \theta}}{\sin \theta} \quad \text{-----} \\ &= \frac{\cos \theta + 1}{\sin \theta} \quad \text{-----} \\ &= \frac{1}{\sin \theta} \\ &= \operatorname{cosec} \theta \end{aligned}$$

- (b) Hence solve the equation  $\cot 3\theta + \frac{\sin 3\theta}{1+\cos 3\theta} = -2$  for  $-90^\circ \leq \theta \leq 90^\circ$ . [4]

$$\operatorname{cosec} 3\theta = -2 \text{-----M1}$$

$$\sin 3\theta = -0.5$$

$$\text{Basic angle} = 30$$

$$3\theta = -30, -150, 210$$

$$\theta = -10, -50, 70 \text{-----}$$

12 (a) Solve  $6^{2x-1} = 4^x \times 5$

[4]

$$\frac{6^{2x}}{6} = 4^x \times 5$$

$$\frac{36^x}{4^x} = 30$$

$$9^x = 30$$

$$x = \log_9 30 \text{ or } 1.548$$

(b) Given that  $\log_x 3 = p$ , express  $\log_3 \frac{27}{x}$  in terms of  $p$ .

[2]

$$\begin{aligned} \log_3 27 - \log_3 x & \text{-----} \\ = 3 - \frac{1}{p} & \text{-----} \end{aligned}$$

(c) Solve the equation  $\lg x - 1 = \lg(x - 1)$


[3]

$$\lg x - \lg 10 = \lg(x - 1)$$

$$\frac{x}{10} = x - 1$$

$$x = \frac{10}{9}$$

- 13 The diagram shows part of the curve  $y = 8 - e^{-3x}$  which crosses the axes at  $A$  and at  $B$ .



- (a) Show that the  $x$  coordinate of  $A$  is  $-\ln 2$ . [3]

$$\begin{aligned} \text{when } y = 0, e^{-3x} &= 8 \dots \\ -3x &= \ln 8 \\ x &= -\frac{1}{3} \ln 8 \dots \\ x &= -\ln 8^{1/3} \\ x &= -\ln 2 \dots \end{aligned}$$

- (b) The normal to the curve at  $B$  meets the  $x$  axis at  $C$ . Find the coordinates of  $C$ . [4]

$$\frac{dy}{dx} = 3e^{-3x} \dots$$

$$\begin{aligned} \text{At } B, x = 0, \text{ so } dy/dx &= 3 \dots \\ \text{Find Gradient of normal at } B &= -1/3 \end{aligned}$$

$$\text{Find } B(0, 7) \dots$$

$$\text{Equation of normal : } y = -\frac{1}{3}x + 7$$

$$\text{So } C(21, 0) \dots$$

- (c) Find the area of the shaded region.

[4]

$$\text{Area of OAB} = \int_{-\ln 2}^0 8 - e^{-3x} dx$$


$$= \left[ 8x + \frac{e^{-3x}}{3} \right]_{-\ln 2}^0$$

$$= 8 \ln 2 - \frac{7}{3}$$

$$\text{Area of triangle OBC} = 0.5 \times 21 \times 7 = 73.5$$

$$\text{Area of shaded region} = 76.7 \text{ units}^2$$

14



The diagram shows a circle, centre  $O$ , with diameter  $AB$ . The point  $C$  lies on the circle. The tangent to the circle at  $A$  meets  $BC$  extended at  $D$ . The tangent to the circle at  $C$  meets the line  $AD$  at  $E$ .

- (a) Prove that triangles  $EOA$  and  $EOC$  are congruent. [3]

$AE = CE$  (tangents from external point are equal)

$AO = OC$  (radius)

$OE = OE$  (common side) -----

Hence triangle  $EOA$  is congruent to triangle  $EOC$  (SSS) -----



(b) Prove that triangles  $ADC$  and  $BAC$  are similar.

[3]

Angle  $ACB = 90$  ( angle in a semi circle )

Angle  $DCA = 90$  ( sum of angles on a straight line )

Hence angle  $ACB =$  angle  $DCA$  ( A ) -----

Angle  $CAD =$  Angle  $CBA$  ( A )  
( angles in alternate segment/ tangent chord theorem ) ----

Hence triangles  $ADC$  and  $BAC$  are similar. ( AA ) -----

(c) Hence prove that  $AC^2 = CD \times BC$ .

[2]

Since they are similar,  $\frac{AC}{BC} = \frac{DC}{AC}$ -----

$AB^2 = BC \times DC$ -----

- End of Paper -

